Diluted Bitumen Affects Multiple Physiological Systems in Sockeye Salmon (*Oncorhynchus nerka*) Embryo to Juvenile Life Stages

Feng Lin,^a Sarah L. Alderman,^b Todd E. Gillis,^b and Christopher J. Kennedy^{a,*}

^aDepartment of Biological Sciences, Simon Fraser University, Burnaby, British Columbia, Canada ^bDepartment of Integrative Biology, University of Guelph, Guelph, Ontario, Canada

Abstract: An understanding of the risks associated with diluted bitumen (dilbit) transport through Pacific salmon habitat necessitates the identification and quantification of hazards posed to early life stages. Sockeye from the embryo to juvenile stage (8 months old) were exposed to four concentrations of the water-soluble fraction of Cold Lake dilbit (summer blend; concentrations of 0, 13.7, 34.7, and 124.5 μ g/L total polycyclic aromatic compounds). Significant mortality (up to 18% over controls) only occurred in the embryo to swim-up fry stage. Impaired growth was seen in the alevin, swim-up, and juvenile stages (maximum reduction 15% in mass but not fork length). Reductions in both critical (maximum 24% reductions) and burst (maximum 47% reductions) swimming speed in swim-up fry and juveniles were seen. Alterations in energy substrate reserves (reductions in soluble protein and glycogen content, elevations in whole-body lipid and triglyceride levels) at all stages may underlie the effects seen in swimming and growth. Dilbit exposure induced a preexercise physiological stress response that affected the recovery of postexercise biochemistry (cortisol, glycogen, lactate, triglyceride concentrations). The transcript abundance of the cytochrome P450 1A gene (*cyp1a*) was quantified in alevin head regions (containing the heart) and in the hearts of swim-up fry and juveniles and showed a concentration-dependent increase in the expression of *cyp1a* at all life stages. *Environ Toxicol Chem* 2022;41:1937–1949. © 2022 SETAC

Keywords: Diluted bitumen; Sockeye salmon; Early life stages; Swimming; Gene expression; Biochemistry

INTRODUCTION

Canada has the world's third largest crude oil reserves, which are estimated at 1.67 trillion barrels, with 96% of proven reserves contained in oil sands deposits located in the Western Canada Sedimentary Basin (Natural Resources Canada, 2020). The extraction of bitumen from the oil sands has increased exponentially over the past decade, with extraction rates projected to increase from the current 2.8 to 4.5 million barrels/day by 2040 (Canada Energy Regulator, 2019). Raw bitumen naturally has high viscosity and density; extracted bitumen is processed and diluted with other, lighter petroleum products (e.g., natural gas condensate or synthetic oil) to facilitate transportation via pipeline (Dew et al., 2015). Diluted bitumen (dilbit; 20%–30% natural gas condensate, 70%–80% bitumen) is the most frequently transported bitumen product in currently employed pipeline networks across North America (Crosby

* Address correspondence to ckennedy@sfu.ca Published online 21 May 2022 in Wiley Online Library (wileyonlinelibrary.com). DOI: 10.1002/etc.5362 et al., 2013; Environment and Climate Change Canada et al., 2013). To cope with increasing global demand for petroleum products, multiple pipeline projects have been proposed in recent years, aiming to increase the exports of Canadian oil sands products (Levy, 2009; National Energy Board [NEB], 2019). The construction of new pipeline and the expansion of existing infrastructure are expected to provide convenient and cost-efficient means for transporting dilbit from remote production sites to coastal regions for refining and eventual overseas shipping (NEB, 2019). The anticipated increase in dilbit transportation (e.g., pipeline, tanker, and rail) raises concerns regarding the potential for a spill event following a pipeline failure or a tanker accident.

Different petroleum products will exhibit unique environmental fate/behaviors and environmental impacts if spilled; limited evidence with dilbit indicates that releases will result in challenging postspill cleanup and habitat recovery (Alsaadi, Hodson, & Langlois, 2018; Dew et al., 2015). For example, it is estimated that nearly 1 million L (up to 30%) of the residual oils from the Kalamazoo River dilbit spill (2010) are associated with sediments and have remained in the river system for years following cleanup efforts (US Environmental Protection Agency, 2013); this suggests that investigations utilizing sublethal and chronic exposure scenarios are needed (Alderman et al., 2018).

Dilbit is a mixture of various petrogenic hydrocarbons (e.g., benzene, toluene, ethylbenzene, xylenes [BTEX], polycyclic aromatic compounds [PACs], and naphthenic acids, among others) with demonstrated toxicity to fish. Acute and sublethal effects following exposure to other crudes or their constituents include developmental defects at early life stages (ELS), impaired growth, reductions in reproductive capacity, changes in behavior, alterations in biochemistry and gene expression, suppressed immune function, genetic damage, and endocrine disruption (Dupuis & Ucan-Marin, 2015; Kennedy, 2014; National Academies of Sciences, Engineering, and Medicine [NASEM], 2016).

Limited studies exist on the toxicity of dilbit to fish species (Dupuis & Ucan-Marin, 2015; NASEM, 2016); therefore, uncertainties regarding the hazards associated with this complex mixture exist. Constituent profiles can vary greatly between products and blends, leading to reservations in predicting dilbit toxicity from data that exist for other crude oils (NASEM, 2016). Early studies on dilbit toxicity focused on embryonic and larval life stages, where effects tend to be greatest in fish exposed to conventional crude oils (see Alsaadi, Madison et al., 2018; Madison, Hodson, & Langlois, 2015; Madison et al., 2017; McDonnell et al., 2019; Philibert et al., 2016). More recent studies investigating effects in older life stages at risk of dilbit exposure support the need for a deeper appreciation of life-specific responses (e.g., fry and juveniles; Alderman et al., 2018, 2020; Alderman, Dindia et al., 2017; Alderman, Lin et al., 2017; Avey et al., 2020; Lin et al., 2020, 2021).

Freshwater and estuarine salmon habitats are at risk because of proposed and existing pipeline and rail routes, as well as the use of marine tanker terminals in the Canadian Pacific Northwest (Levy, 2009; Raincoast Conservation Foundation, 2018). As one of the most productive salmon migration routes in the world, the Fraser River watershed and its estuary serve as vital spawning and nursery habitat for all five species of Pacific salmon (Henderson & Graham, 1998; Labelle, 2009). There is an increasing body of evidence suggesting that dilbit exposure can negatively affect the survival, early development, and critical physiological systems of Pacific salmon at extremely low environmental concentrations. These effects include delayed hatching time, mortality during embryonic development, deformities, impairment of growth, and alteration of body composition (Alderman et al., 2018). Older ELS are also affected by dilbit exposure. For example, exposed 1+-year-old sockeye exhibit altered gene expression, decreased swimming ability, as well as alterations in cardiac tissues and the plasma proteome (Alderman, Dindia et al., 2017; Alderman, Lin et al., 2017). The potential for longer-term exposures to dilbit following a spill in salmon habitat and the growing evidence of pronounced effects in salmon were the impetus for the present study. In the present study, the effects of a chronic dilbit exposure on developing sockeye through several life stages from the embryo to the juvenile stage were investigated using a suite of endpoints known to have direct relevance to salmon

survival and performance (survival, growth, biochemistry, gene expression, swim performance, and exercise recovery).

MATERIALS AND METHODS

Fish

Sockeye gametes were obtained from the Upper Pitt River Hatchery (Fisheries and Oceans Canada) and fertilized according to standard procedures (Ontario Ministry of Natural Resources, 2009). Embryos were incubated in heath trays (MariSource; 372 embryos per tray; mean mass 0.24 ± 0.11 g [mean ± standard deviation]) supplied with dechlorinated municipal water (flow rate 6 L/min; dissolved $O_2 > 95\%$ saturation, hardness $6.12 \text{ mg/L} \text{ CaCO}_3$, dissolved organic carbon < 1 mg/L, pH 7.0) at 11.3 °C in the dark until the swim-up fry stage (no visible external yolk sac). Mortality was recorded daily under red light, and dead embryos were immediately removed from trays. Swim-up fry were collected from rearing trays and transferred to 250-L fiberglass tanks supplied with dechlorinated water at 13 °C (flow rate 7.5 L/min and 12:12-h light:dark photoperiod). Fry were fed 5% body weight/day commercial salmonid feed (Skretting Canada), which was increased weekly according to a growth equation that included a feed conversion efficiency of 20% (Meador et al., 2006) until fish were 8 months of age (juveniles). The care and use of all fish were approved by the University Animal Care Committee at Simon Fraser University following Canadian Council on Animal Care quidelines (1315B-20).

Exposure

The water-soluble fraction of dilbit was generated as previously described (Alderman, Dindia et al., 2017; Kennedy & Farrell, 2005). In brief, Siproax[®] ceramic beads (Aquatic Eco-Systems) were soaked (except controls) in unweathered Cold Lake Blend summer dilbit (COOGER, DFO) for 24 h and then placed into polyvinyl chloride columns (16 cm diameter x 80 cm length) supplied with an upward-directed continuous flow of dechlorinated municipal water (6 L/min). Varying the number of beads/column provided four different water-soluble fraction concentrations (in duplicate); columns were "recharged" every 14 days with newly soaked beads. Water containing the water-soluble fraction of dilbit flowed into 500-L fiberglass header tanks and was then distributed into heath stacks. Embryos were exposed to dilbit in duplicate heath stacks immediately postfertilization and continued until fish reached the swim-up stage. At this stage, fish were transferred into 200-L fiberglass tanks (n = 200 fish in each duplicate tank) supplied with dilbit-water-soluble fraction water and exposed for a further 90 days until fish were 8 months of age (8-month total exposure). The detailed experimental design is depicted in Figure 1.

Water samples were collected from duplicate header tanks at 0, 7, and 14 days after initiation of water-soluble fraction generation and analyzed for individual PACs using gas chromatography-mass spectrometry, as previously described



Sockeye salmon eggs were fertilized

Exposure started 2 dpf

Developing

embryo

2 dpf

FIGURE 1: Schematic of the experimental design. ELS = early life stage; WSF = water-soluble fraction; dpf = days postfertilization.

(Alderman, Dindia et al., 2017). Individual PAC concentrations were measured by Axys Analytical Services following standard procedures. All samples were spiked with deuterated surrogate standards prior to dichloromethane extraction and cleanup with silica column chromatography. Low-resolution mass spectrometry using an RTX-5 capillary gas chromatography column operated in the electron impact ionization mode using multiple ion detection was used, acquiring at least one characteristic ion for each target analyte and surrogate standard. Reporting limits for individual compounds ranged from 0.13 to 1.13 ng/L (mean 0.42 ng/L), and the mean percentage recovery was 100.3%. Concentrations of target PACs were calculated using the isotope dilution method of quantification and expressed as percentage of total PAH (Alderman, Dindia et al., 2017).

Tissue collection

Developing alevins (n = 35-94 days postfertilization [dpf]), swim-up fry (n = 44-147 dpf), and juveniles (n = 44-237 dpf)were randomly sampled from each exposure group (and replicate tanks), euthanized with buffered tricaine mesylate (MS-222; 1 g/L), and weighed and measured for length. For biochemical analysis, a subset of these fish (n = 16-20)treatment) were frozen in liquid N_2 and then transferred to -80 °C until analysis for whole-body lipid and triglyceride content. For gene transcript analysis, the head region of alevins (bisected at the rostral boundary of the yolk sac and perpendicular to the body axis) or the isolated hearts (fry and juveniles) were individually frozen in liquid N_2 and stored at -80 °C for reverse-transcription quantitative polymerase chain reaction (RT-qPCR) analysis (n = 8/developmental stage, n = 8/treatment group, n = 4/replicate tank).

Swim tests

Critical swimming speed (U_{crit}) and burst swimming speed (U_{burst}) tests (Farrell, 2008; Osachoff et al., 2014) were performed for swim-up fry and juveniles (n = 10 from each treatment) using an isolated/blacked-out mini-swim tunnel system (Loligo[®] Systems). The swim tunnel system (a 1.5-L cylindrical glass chamber submerged inside a reservoir) was temperatureregulated by a custom chilled bath circulator, and dissolved oxygen was maintained at >95% by constant aeration. Fish were first acclimated for 20 min at a water velocity of 1.5 body lengths per second (BL/s) before the swim tests. Water velocity was then increased by 1.5 BL/s every 20 min until fish were exhausted (Farrell, 2008). The U_{burst} test was initiated at a water velocity of 1.0 BL/s (Farrell, 2008; Nendick et al., 2009), rapidly increased to 2 BL/s over a 1-min interval, and subsequently increased by 0.5 BL/s every 1 min. Both tests were complete when exhausted fish were inactive on the rear baffle for over 2 s and would not resume swimming after a brief decrease in water velocity. Fish were immediately removed from the tunnel and euthanized in buffered MS-222, and wet weight (grams) and fork length (centimeters) were recorded. Values for $U_{\rm crit}$ and U_{burst} were calculated according to Farrell (2008). The cross-sectional area of all swim-tested fish was found to be <10% of swim tunnel cross-sectional area, and fish density was <0.2 g/L; therefore, U_{crit} and U_{burst} values were not corrected for solid blocking effects (Bell & Terhune, 1970). Euthanized U_{crit}-tested fish were frozen in liquid nitrogen and immediately transferred to -80 °C for postexercise body composition analysis, as described below (see Body composition).

Biochemical measurements

Whole-body lipid content was measured using a standard protocol (Folch et al., 1957). Preweighed fish were thawed on ice and minced into pieces in envelopes (Whatman filter paper) that were then sealed and saturated in a chloroform:methanol (2:1) mixture (solvent:tissue 20:1), incubated for 20 min in a glass container, and shaken at 30 rpm. Samples were then washed with chloroform and dried in an oven (60 °C) for 24 h. Total-body lipid content of each individual fish was calculated by subtracting the sample's original wet weight by the net weight of the dried sample.

Total soluble protein content was quantified in preweighed whole fish that were thawed on ice and homogenized in nine volumes of lysis buffer (0.5 M Tris-HCl and 0.1 mM ethylenediaminetetraacetic acid at pH 8) using a Mixer Mill homogenizer (model MM 300; Qiagen; Cassidy et al., 2016). Crude homogenates were centrifuged at 13,000g at 4 °C for 60 min, and the soluble protein content in supernatants was

measured using a Bradford protein assay kit with bovine serum albumin as the standard (catalog no. 5000002; Bio-Rad).

For whole-body pre- and postexercise cortisol, glycogen, triglyceride, and lactate concentrations, preweighed alevin or fry were thawed on ice and homogenized in 0.2 M sodium citrate buffer at pH 5 (catalog no. 200-675-3; EMD Chemicals) using a Tissue Tearor (Fisher Scientific). Each crude homogenate sample was aliquoted into separate microcentrifuge tubes and stored at -80 °C until subsequent analysis. The glycogen concentration was determined following a standard protocol (Weber et al., 2008) using Type IX bovine liver glycogen as a standard (C940M53; Sigma-Aldrich). The whole-body triglyceride concentration was determined following the microplate spectrophotometric assay protocol as described in Weber et al. (2003). Whole-body lactate content of each fish was determined using a commercial colorimetric assay kit (catalog no. 120001400P; Eton Bioscience) performed according to the manufacturer's protocol. Total cortisol concentration was quantified using a commercial enzyme-linked immunosorbent assay kit (catalog no. EA65; Oxford Biomedical Research; McPhee & Janz, 2014). All colorimetric assays were performed in duplicate using an Epoch[™] 2 microplate spectrophotometer (Bio-Tek) and a Corning[®] 96-well microplate (Greiner Bio-One International).

RT-qPCR analysis

Transcript abundances of cytochrome P450 1a (*cyp1a*; Phase I biotransformation) and ribosomal protein L8 (*rpl8*; housekeeping gene) were quantified in alevin head regions (containing the heart) and in the hearts of swim-up fry and juveniles following standard quality control guidelines (Bustin et al., 2009) and using total RNA extraction, complementary DNA synthesis, and RT-qPCR methods exactly as previously described (Alderman et al., 2018). Primer sequences were *cyp1a* (F: tcatcaacgacggcaaga, R: gttcaccaagcccaacag, 110% efficiency) and *rpl8* (F: ttggtaatgttctgcctgtg, R: gggttgtggga-gatgactg, 103% efficiency). Data were normalized to the abundance of the stably expressed reference gene, *rpl8*.

Statistical analysis

No statistically significant tank effect was found (using either one-factor or two-factor analysis of variance [ANOVA]) in which *tank* was included as a random factor); therefore, data from replicate tanks were pooled for all analyses. Mortality, body composition measures (including condition factor [mass/length³]), and swim test data were analyzed using one-factor ANOVA and Tukey's multiple comparison test ($\alpha = 0.05$). The preexercise and postexercise biochemical data from different treatment groups were combined and compared using two-factor ANOVA followed by Tukey's multiple comparison test ($\alpha = 0.05$). Differences in transcript abundances were determined using one-factor ANOVA and Tukey's multiple comparisons test, and the relative abundance of *cyp1a* in the hearts of swim-up fry and 8-month-old juveniles was compared with a

two-factor ANOVA and Sidak's multiple comparisons test (n = 8; $\alpha = 0.05$).

RESULTS

Water chemistry

Water samples collected (two composite samples per concentration) at 0 days confirm the presence of PAC in experimental tanks supplied with water-soluble fraction of dilbit, with initial total PAC (TPAC) values ranging from $0.2 \,\mu$ g/L (control) to $13.7 \,\mu$ g/L (low) to $34.7 \,\mu$ g/L (medium) to $124.5 \,\mu$ g/L (high). The initial TPAC concentrations in water-soluble fraction of dilbit exposures are used hereafter to

(A) Embryo to swim-up stage



Treatment

FIGURE 2: Cumulative mortality of embryos exposed to dilbit. The initial total polycyclic aromatic compound concentrations for each water-soluble fraction were 0.2 (control), 13.7 (low), 34.7 (medium), and 124.5 (high) μ g/L. Time of exposure was (**A**) 147 days for embryo to swim-up fry and (**B**) 90 days for swim-up stage to 8-month-old juveniles. One-factor analysis of variance and Tukey's multiple comparison test were used to test for significant differences between water-soluble fraction treatment groups. Bars that do not share a common letter are statistically different (p < 0.05).

Life stage	Treatment	Body mass (mg)	Fork length (mm)	Condition factor	n
Alevin	Control	182.9 ± 2.0A	22.6 ± 0.2A	1.61 ± 0.03A,B	70
	Low	184.1 ± 2.1A	22.2 ± 0.1A	1.70 ± 0.03A	70
	Medium	178.9 ± 1.9A,B	22.4 ± 0.1A	$1.60 \pm 0.02B$	70
	High	173.3 ± 2.4B	22.3 ± 0.2A	1.58 ± 0.03B	70
Swim-up fry	Control	161.9 ± 0.9A	28.8 ± 0.1A	0.68 ± 0.003B	227
	Low	163.7 ± 0.7A	28.4 ± 0.1B	0.72 ± 0.003A	179
	Medium	140.2 ± 0.9B	27.0 ± 0.1C	0.72 ± 0.004A	177
	High	137.4 ± 1.6B	26.9 ± 0.1C	0.71 ± 0.006A	92
Juveniles	Control	673.8 ± 6.9A	39.7 ± 0.2A	1.08 ± 0.008A	368
	Low	651.3 ± 6.9A	39.3 ± 0.2A	1.08 ± 0.008A	373
	Medium	561.5 ± 6.8B	39.4 ± 0.2A	0.92 ± 0.009B	367
	High	583.8 ± 7.1B	39.6 ± 0.2A	0.94 ± 0.008B	363

TABLE 1: Wet weight, fork length, and Fulton's condition factor of fish exposed to water-soluble fraction of Cold Lake Blend dilbit

The initial total polycyclic aromatic compound concentrations for each water-soluble fraction were 0.2 (control), 13.7 (low), 34.7 (medium), and 124.5 (high) μ g/L. Time of exposure was 147 days for embryo to swim-up fry and 90 days for swim-up stage to 8-month-old juveniles. Data are means ± standard error for n = 20 fish. Treatments that do not share a common letter are statistically different (p < 0.05).



FIGURE 3: Cytochrome P450 1a (*cyp1a*) expression in (**A**) alevin head regions (50% yolk) and isolated whole hearts of (**B**) swim-up fry and (**C**) 8-month-old juveniles exposed to various concentrations of dilbit. The initial total polycyclic aromatic compound concentrations for each watersoluble fraction were 0.2 (control), 13.7 (low), 34.7 (medium), and 124.5 (high) μ g/L. Expression was normalized to the housekeeping gene *rpl8*. Within each life stage, data are expressed as fold-change from unexposed controls, and boxes that do not share a common letter are significantly different (one-way analysis of variance [ANOVA] and Tukey's test, n = 8; p < 0.001). The mean abundance of *cyp1a* in swim-up fry versus juveniles at each concentration is shown in (**D**), with significant differences between life stages indicated with an asterisk (two-way ANOVA and Sidak's post hoc test, n = 8; $p_{\text{interaction}} = 0.013$). *rpl8* = ribosomal protein L8.



FIGURE 4: Whole-body total (**A**) lipid, (**B**) triglyceride, and (**C**) protein content in alevins exposed to dilbit. The initial total polycyclic aromatic compound concentrations for each water-soluble fraction were 0.2 (control), 13.7 (low), 34.7 (medium), and 124.5 (high) μ g/L. Within each plot, + indicates mean for n = 20 fish; boxes that do not share a common letter are statistically different (p < 0.05).

designate treatment groups. In this system, TPAC concentrations decreased by 50%–70% between 0 and 12 days and then changed very little between 12 and 25 days (Alderman, Dindia et al., 2017; Lin et al., 2020). Component breakdown for TPAC shows that smaller and more volatile hydrocarbons (e.g., naphthalenes) predominate initially, with larger PACs (e.g., phenanthrenes) becoming relatively more abundant with time (Lin et al., 2020).

Mortality and growth

Cumulative mortalities in embryos and swim-up fry in the medium and high water-soluble fraction treatments were significantly higher (medium vs. control, p = 0.0482; high vs. control, p < 0.01) than in controls; however, no significant mortality occurred in older fish from exposure (p > 0.05; Figure 2). Body mass was significantly lower in alevins exposed to the highest concentration compared with controls, and at the swim-up stage significantly lower body mass was seen in both the medium and high treatment groups (medium and high both p < 0.01), a trend which continued to the juvenile stage (Table 1). Significantly reduced length was only seen in water-soluble fraction-exposed swim-up fry (Table 1). Condition factor was marginally but significantly higher in swim-up fry and lower in juveniles at the higher two concentrations compared with controls (Table 1).

Molecular responses

There was a concentration-dependent increase in transcript abundance of *cyp1a* at all stages examined, with maximal responses of 72-fold (50% yolk sac; Figure 3A), 9-fold (swim-up fry; Figure 3B), and 162-fold (8-month-old juveniles; Figure 3C) relative to unexposed controls. The apparent low response in swim-up fry, however, is relative to an already high background expression of *cyp1a*. Specifically, expression of *cyp1a* in the hearts of unexposed control swim-up fry was 32-fold greater than in the hearts of unexposed control juveniles, and the peak response observed in fish exposed to the highest concentration of dilbit was also relatively higher in swim-up fry hearts (Figure 3D).

Body composition

Developing alevins exposed to dilbit exhibited increased total-body lipid and triglyceride concentrations and a reduced total soluble protein content compared with controls (Figure 4); whole-body lipid contents in the medium and high treatment groups were 1.9-fold and 2.2-fold higher than in controls (p=0.048, p=0.015), and triglyceride levels in these two groups were elevated by 1.6-fold and 2.0-fold, respectively (p=0.034, p<0.01). Lower protein content (35%, high vs. control, p = 0.025; 27%, medium vs. control, p < 0.01, respectively) was also seen. Swim-up fry in the two higher treatment groups exhibited an increased total lipid content, 2.0-fold and 2.2-fold higher than controls (Figure 5; medium vs. control, p < 0.01; high vs. control, p < 0.01) as well as lower whole-body soluble protein levels (medium vs. control, by 33.6%, p = 0.0241; high vs. control, 42.1%, p < 0.01; Figure 5). Juveniles in the high exposure group had a higher lipid content (1.7-fold, p < 0.01) and lower whole-body soluble protein levels



FIGURE 5: Total lipid content (**A**,**B**) and total protein content in (**C**,**D**) swim-up fry and juveniles exposed to dilbit. The initial total polycyclic aromatic compound concentrations for each water-soluble fraction were 0.2 (control), 13.7 (low), 34.7 (medium), and 124.5 (high) μ g/L. Within each plot, + indicates mean for n = 16 fish; boxes that do not share a common letter are statistically different (p < 0.05).

(by 27%) in the high treatment group (p = 0.048; Figure 5) compared with controls.

Swim performance

Values of $U_{\rm crit}$ and $U_{\rm burst}$ were affected by exposure in both swim-ups and juveniles. Values of $U_{\rm crit}$ for swim-up fry in the medium and high treatments were 20% lower than those of controls (Figure 6; medium vs. control, p < 0.01; high vs. control, p < 0.01; for juveniles, $U_{\rm crit}$ was reduced by 22.8% compared with controls in the high treatment group (p < 0.01). Values of $U_{\rm burst}$ were decreased in swim-up fry in the medium and high treatments (26% and 39%, respectively) compared with controls (Figure 6; medium vs. control, p < 0.01; high vs. control, p < 0.01). For the juveniles, the medium and high groups exhibited decreased $U_{\rm burst}$ by 16.3% and 22.2%, respectively, compared with controls (p < 0.01).

Pre- and postexercise biochemistry

Preexercise (baseline) whole-body cortisol concentrations in swim-ups and juveniles were higher (range 1.7- to 2.7-fold) in

fish exposed to the medium and high treatment groups compared with controls (p < 0.01, p = 0.037; Figure 7). Fish that underwent the $U_{\rm crit}$ test exhibited a significant elevation in whole-body cortisol concentrations in controls (2.1-fold, p = 0.049) and the low treatment group (2.2-fold, p = 0.023) in swim-up fry. In contrast, postexercise cortisol concentrations in fish from the medium and high treatments were not significantly different from preexercise baseline values. This lack of an exercise-induced cortisol increase was consistently observed in juveniles in all three dilbit treatment groups (Figure 7).

Preexercise body glycogen reserves were lower in fish in the higher treatment groups compared with controls (medium vs. control, p < 0.01; high vs. control, p < 0.01; Figure 7). The $U_{\rm crit}$ test resulted in significant reductions in total-body glycogen content in both control and water-soluble fraction-exposed swim-ups and juveniles; however, in juveniles, exposure to the highest concentration further reduced glycogen stores compared with controls (p < 0.01; Figure 7).

For both swim-ups and juveniles, preexercise lactate levels of exposed fish were not significantly different from those of controls (p > 0.05; Figure 8). For both swim-up fry and juveniles, significant increases in whole-body lactate levels were



FIGURE 6: Critical and burst swimming speed of (**A**,**B**) swim-up fry and (**C**,**D**) juveniles exposed to dilbit. The initial total polycyclic aromatic compound concentrations for each water-soluble fraction were 0.2 (control), 13.7 (low), 34.7 (medium), and 124.5 (high) μ g/L. Within each plot, **+** indicates mean for n = 10 fish; boxes that do not share a common letter are statistically different (p < 0.05). $U_{crit} =$ critical swimming speed; $U_{burst} =$ burst swimming speed; BL = body length.

observed following the U_{crit} trial in both control and exposed fish (p < 0.01; Figure 8). For both swim-up fry and juveniles, only fish exposed to the highest water-soluble fraction had higher postexercise lactate when compared with the controls (p = 0.0225, p = 0.0193).

Whole-body triglyceride levels in preexercised swim-ups were significantly higher in the medium and high treatment groups (p < 0.01 and p < 0.01, respectively) and in the high treatment group in juveniles compared with controls (p < 0.01, Figure 8). Exercise caused significant decreases in control and low treatment swim-ups (p = 0.0189, p = 0.025) and in juveniles (p = 0.0484, p = 0.0275). For both swim-up fry and juveniles, fish exposed to the highest concentration of water-soluble fraction did not exhibit a significant decrease in whole-body triglyceride following exhaustive swimming exercise as was seen in other groups (p > 0.05; Figure 8).

DISCUSSION

1944

Evidence is growing that Pacific salmon ELS, particularly developing embryos and alevins, are more sensitive to dilbit

than other species and that tolerance in salmon increases with age. Dilbit-induced lethality in developing sockeye was life stage-dependent; embryos through the swim-up stage were most sensitive with mortality occurring at TPAC as low as 34.7 µg/L. Alderman et al. (2018) reported higher sensitivity, with mortality in sockeye embryos at TPAC concentrations as low as $4 \mu g/L$; however, at a similar concentration (35 $\mu g/L$ TPAC), overall mortality was similar (8% vs. 13%). No mortality was seen in dilbit-exposed 1+-year sockeye parr exposed to concentrations between 3.5 and 66.7 µg/L (Alderman, Dindia et al., 2017). Developing pink salmon (Oncorhynchus gorbuscha) embryos exhibited mortality at a similar total polycyclic aromatic hydrocarbon (TPAH) range (18-48 µg/L) using a similar dosing method (Heintz et al., 1999). In contrast, larval fathead minnow (Pimephales promelas) and inland silverside (Menidia beryllina) exhibited no acute lethality at TPAH of 8-40 µg/L (Barron et al., 2018). Low lethal toxicity has also been reported in zebrafish (Danio rerio, 28 µg/L), fathead minnow, Japanese medaka (Oryzias latipes), and yellow perch (Perca flavescens; at TPAH <100 µg/L) during embryonic development (Alsaadi, Madison et al., 2018; Madison, Hodson, & Langlois, 2015;



FIGURE 7: Preexercise and postexercise whole-body cortisol content and glycogen content in (**A**,**B**) swim-up fry and (**C**,**D**) juveniles exposed to dilbit. The initial total polycyclic aromatic compound concentrations for each water-soluble fraction were 0.2 (control), 13.7 (low), 34.7 (medium), and 124.5 (high) μ g/L. Within each plot, + indicates mean for n = 10 fish; boxes that do not share a common letter are statistically different (p < 0.05).

Madison et al., 2017; McDonnell et al., 2019; Philibert et al., 2016).

Body mass, but not overall length, was reduced in sockeye at all life stages when exposed to dilbit; and marginal inconsistent effects were seen in condition factor. Similar results were reported for various petroleum products in other salmonid species (Atlantic salmon, *Salmo salar* [Vignier et al., 1992]; cutthroat trout, *Oncorhynchus clarkii* [Woodward et al., 1983]; rainbow trout, *Oncorhynchus mykiss* [Lockhart et al., 1996]; pink salmon [Wang et al., 1993]; and Chinook salmon, *Oncorhynchus tshawytscha* [Meador et al., 2006]). Multiple mechanisms have been suggested to underlie growth reductions, including the suppression of feeding behavior and decreases in food conversion efficiency (Moles & Rice, 1983; Vignier et al., 1992), physiological stress (Kennedy & Farrell, 2005, 2006; Kochhann et al., 2015), and elevations in metabolic rate (dos Santos et al., 2006; Klinger et al., 2015).

The availability and utilization of critical energy substrates have direct bearing on the success of embryonic development and the growth of fish (Srivastava & Brown, 1991). Dilbit-induced alterations in whole-body lipid, triglyceride, glycogen, and protein concentrations at all stages suggest that disturbances in lipid and protein metabolism or their utilization likely underlie the alterations in mass, length, and calculated condition factor seen at the whole-animal level. Altered lipid stores and reductions in free protein levels have been reported in salmon alevins exposed to dilbit and are suggested to underlie delayed development and shortened body lengths (Alderman et al., 2018). Similarly, the transient exposure of polar cod (Boreogadus saida) to North Slope crude oil during embryogenesis caused significant elevations in lipid content (e.g., triacylglycerols, free fatty acids, sterols), reductions in posthatching body size, and poor survival (Laurel et al., 2019). In developing Atlantic haddock (Melanogrammus aeglefinus), crude oil exposure disrupted yolk lipid utilization and the biosynthesis of intrinsic cholesterol (Sørhus et al., 2017). Disruptions of lipid utilization may result in the use of protein and carbohydrate as alternative substrates, affecting their concentrations. Increased oxygen consumption in developing mahi-mahi (Coryphaena hippurus) exposed to Deep water Horizon crude oil is possibly fueled by enhanced endogenous protein catabolism (Pasparakis et al., 2016). Transcriptomic studies in larval mahi-mahi and red drum (Sciaenops ocellatus) have shown that pathways involved in amino acid metabolism and protein digestion are significantly altered following exposure to Deep water Horizon crude oil (Xu et al., 2016, 2017).

As in the present study, effects on swimming in various teleosts can occur at concentrations of TPAC as low as $0.23-200 \mu g/L$ (Hicken et al., 2011; Johansen & Esbaugh, 2017; Kennedy & Farrell, 2006; Mager et al., 2014). Aberrant molecular responses, functional deficits, and morphological/histopathological alterations during cardiogenesis (Alsaadi, Madison, et al., 2018; Madison, Hodson et al., 2015; Madison et al., 2017; McDonnell et al., 2019; see also Brette et al., 2014; Incardona, 2017); remodeling in cardiac tissues



FIGURE 8: Preexercise and postexercise whole-body lactate and triglyceride content in (**A**,**B**) swim-up fry and (**C**,**D**) juveniles exposed to dilbit. The initial total polycyclic aromatic compound concentrations for each water-soluble fraction were 0.2 (control), 13.7 (low), 34.7 (medium), and 124.5 (high) μ g/L. Within each plot, + indicates mean for n = 10 fish; boxes that do not share a common letter are statistically different (p < 0.05).

(Alderman, Dindia et al., 2017; Alderman, Lin et al. 2017); and disruptions to cardiovascular capacity (Johansen & Esbaugh, 2017; Nelson et al., 2017; Stieglitz et al., 2016) are often linked to impaired swimming performance in fish exposed to dilbit. In the present study, an induction of *cyp1a* in the heart was observed in dilbit-exposed swim-up fry and juveniles, consistent with cardiotoxicity as a driving mechanism for the observed reductions in swimming performance.

Routine, sustained, and prolonged swimming are primarily fueled by triglyceride oxidative metabolism in slow-twitch red skeletal muscle; and sprint/burst swimming is fueled by glycolytic metabolism in fast-twitch white muscle (Hammer, 1995; Moyes & West, 1995). Elevations in whole-body triglycerides in exposed fish did not provide advantages in swimming performance, and unchanged triglyceride content post-Ucrit test may reflect a decreased lipolytic capacity; a reduced lipid utilization during the aerobic exercise and diminished carbohydrate availability for anaerobic bursting may underlie impairment. Similarly, Avey et al. (2020) reported that $U_{\rm crit}$ was not affected in Atlantic salmon smolts exposed to watersoluble fraction of dilbit (up to 67.9 µg/L TPAC), but the fish exhibited a reduced reliance on lipid metabolism for adenosine triphosphate in the heart. Exposure to a lower water-soluble fraction concentration at 9.65 µg/L TPAC resulted in an increased reliance on anaerobic metabolism in both cardiac and red skeletal muscle (Avey et al., 2020). Burst swimming is almost exclusively fueled anaerobically through the utilization of muscle glycogen, and reductions in $U_{\rm burst}$ may be directly attributed to the lowered body glycogen stores seen prior to the swim trial. A greater accumulation of lactate and depletion of glycogen postexercise in fish exposed to dilbit suggest a potentially enhanced anaerobic debt during burst swimming.

Dilbit exposure activated a physiological stress response, followed by a short-term hyperglycemic response similar to the response induced by crude oil exposure; this has been attributed to the irritant properties of the lighter, more volatile, and acutely toxic components of oil (e.g., naphthalenes, BTEX, and naphthenic acids; Kennedy & Farrell, 2005, 2006; P. Thomas et al., 1980). Cortisol elevation can reduce feeding and food conversion efficiency (Gregory & Wood, 1999; Madison, Tavakoli, et al., 2015) and may be responsible for the reduced growth seen in the present study. Consistent with cortisol's key role in mediating the peripheral mobilization of energetic substrates during stress, stress may have resulted in the catabolism of body carbohydrate and protein (Milligan, 2003; Mommsen et al., 1999). However, elevated cortisol generally increases peripheral and hepatic lipolysis through increases in lipase activity (Baltzegar et al., 2014), increased glycerol utilization (Vijayan et al., 1991), and reductions in hepatic lipogenic potential (López-Patiño et al., 2014), which is not consistent with the higher lipid and triglyceride content seen in exposed fish that were stressed.

Exhaustive exercise resulted in increased circulating cortisol concentrations, which did not increase over preexercise levels in

response to exercise, data that are contrary to those seen in other species acutely exposed to crude oil (Kennedy & Farrell, 2006; R. E. Thomas & Rice, 1987). Chronic exposure, however, can cause a muted cortisol response following exercise (Kennedy & Farrell, 2005, 2006). The main role of the cortisol-induced stress response is to supply an immediate energy source for fuelintensive behaviors and physiological processes, and a deemphasized cortisol response can be considered maladaptive. Repeated pulse exposures to petroleum may cause hyperactivity and exhaustion of cortisol-producing cells (Hontela, 1997); act as an endocrine disruptor, targeting pituitary or adrenocortical tissues (Dorval et al., 2003) and affecting multiple sites in the hypothalamic–pituitary–interrenal axis (Kennedy & Farrell, 2005); or result in the necrosis of interrenal tissues (DiMichele & Taylor, 1978).

CONCLUSIONS

Chronic exposure of sockeye to dilbit significantly reduced survival and growth, impaired aerobic and anaerobic swimming performance, and altered body biochemical composition as well as cardiac gene expression, providing evidence that this complex mixture likely has multiple targets, resulting in a complex suite of toxicological outcomes. Dilbit release into the natural habitat of Pacific salmon, under similar exposure scenarios, is likely to produce adverse effects that will affect the viability and sustainability of local salmon populations.

Acknowledgments—We thank the Kennedy Lab technician, G. Su, for experimental support and biochemical analysis. We thank the Simon Fraser University animal care staff and B. Leighton for extensive fish care and facility support. The present study was supported by grants from the National Contaminants Advisory Group at Fisheries and Oceans Canada (R529320; to C. J. Kennedy).

Conflict of Interest—The authors declare no conflict of interest.

Author Contributions Statement—Christopher J. Kennedy: Funding acquisition; Conceptualization; Project administration; Resources; Writing—review & editing. Todd E. Gillis: Funding acquisition; Conceptualization; Project administration; Resources; Writing—review & editing. Feng Lin: Investigation; Writing—original draft; Writing—review & editing. Sarah L. Alderman: Investigation; Writing—review & editing.

Data Availability Statement—Data, associated metadata, and calculation tools are available from the corresponding author (ckennedy@sfu.ca).

REFERENCES

Alderman, S. L., Dilkumar, C. M., Avey, S. R., Farrell, A. P., Kennedy, C. J., & Gillis, T. E. (2020). Effects of diluted bitumen exposure and recovery on the seawater acclimation response of Atlantic salmon smolts. *Aquatic Toxicology*, 221, Article 105419. https://doi.org/10.1016/j.aquatox.2020. 105419

- Alderman, S. L., Dindia, L. A., Kennedy, C. J., Farrell, A. P., & Gillis, T. E. (2017). Proteomic analysis of sockeye salmon serum as a tool for biomarker discovery and new insight into the sublethal toxicity of diluted bitumen. *Comparative Biochemistry and Physiology Part D: Genomics and Proteomics*, 22, 157–166. https://doi.org/10.1016/j.cbd. 2017.04.003
- Alderman, S. L., Lin, F., Farrell, A. P., Kennedy, C. J., & Gillis, T. E. (2017). Effects of diluted bitumen exposure on juvenile sockeye salmon: From cells to performance. *Environmental Toxicology and Chemistry*, 36, 354–360. https://doi.org/10.1002/etc.3533
- Alderman, S. L., Lin, F., Gillis, T. E., Farrell, A. P., & Kennedy, C. J. (2018). Developmental and latent effects of diluted bitumen exposure on early life stages of sockeye salmon (*Oncorhynchus nerka*). Aquatic Toxicology, 202, 6–15. https://doi.org/10.1016/j.aquatox.2018.06.014
- Alsaadi, F., Hodson, P. V., & Langlois, V. S. (2018). An embryonic field of study: The aquatic fate and toxicity of diluted bitumen. *Bulletin of Environmental Contamination and Toxicology*, 100, 8–13. https://doi. org/10.1007/s00128-017-2239-7
- Alsaadi, F. M., Madison, B. N., Brown, R. S., Hodson, P. V., & Langlois, V. S. (2018). Morphological and molecular effects of two diluted bitumens on developing fathead minnow (*Pimephales promelas*). Aquatic Toxicology, 204, 107–116. https://doi.org/10.1016/j.aquatox.2018.09.003
- Avey, S. R., Kennedy, C. J., Farrell, A. P., Gillis, T. E., & Alderman, S. L. (2020). Effects of diluted bitumen exposure on Atlantic salmon smolts: Molecular and metabolic responses in relation to swimming performance. *Aquatic Toxicology*, 221, Article 105423. https://doi.org/10.1016/j. aquatox.2020.105423
- Baltzegar, D. A., Reading, B. J., Douros, J. D., & Borski, R. J. (2014). Role for leptin in promoting glucose mobilization during acute hyperosmotic stress in teleost fishes. *Journal of Endocrinology*, 220, 61–72. https://doi. org/10.1530/JOE-13-0292
- Barron, M. G., Conmy, R. N., Holder, E. L., Meyer, P., Wilson, G. J., Principe, V. E., & Willming, M. M. (2018). Toxicity of Cold Lake Blend and Western Canadian Select dilbits to standard aquatic test species. *Chemosphere*, 191, 1–6. https://doi.org/10.1016/j.chemosphere.2017.10.014
- Bell, W. H., & Terhune, L. D. B. (1970). Water tunnel design for fisheries research (Technical Report 195). Fisheries Research Board of Canada.
- Brette, F., Machado, B., Cros, C., Incardona, J. P., Scholz, N. L., & Block, B. A. (2014). Crude oil impairs cardiac excitation–contraction coupling in fish. *Science*, 343, 772–776. https://doi.org/10.1126/science.1242747
- Bustin, S. A., Benes, V., Garson, J. A., Hellemans, J., Huggett, J., Kubista, M., Mueller, R., Nolan, T., Pfaffl, M. W., Shipley, G. L., Vandesompele, J., & Wittwer, C.T. (2009). The MIQE guidelines: Minimum information for publication of quantitative real-time PCR experiments. Clinical Chemistry, 55, 611–622. https://doi.org/10.1373/clinchem.2008.112797
- Canada Energy Regulator. (2019). Canada's energy futures 2018 supplement: Oil sands production. https://www.cer-rec.gc.ca/nrg/ntgrtd/ftr/ 2018lsnds/index-eng.html
- Cassidy, A. A., Saulnier, R. J., & Lamarre, S. G. (2016). Adjustments of protein metabolism in fasting Arctic charr, *Salvelinus alpinus*. *PLoS One*, 11, Article e0153364. https://doi.org/10.1371/journal.pone.0153364
- Crosby, S., Fay, R., Groark, C., Kani, A., Smith, J. R., Sullivan, T., & Pavia, R. (2013). Transporting Alberta oil sands products: Defining the issues and assessing the risks (NOAA Technical Memorandum No. NOS OR&R 44). US Department of Commerce, National Oceanic and Atmospheric Administration, National Ocean Service. https://repository.library.noaa.gov/ view/noaa/2670/noaa_2670_DS1.pdf
- Dew, W. A., Hontela, A., Rood, S. B., & Pyle, G. G. (2015). Biological effects and toxicity of diluted bitumen and its constituents in freshwater systems. *Journal of Applied Toxicology*, 35, 1219–1227. https://doi.org/10. 1002/jat.3196
- DiMichele, L., & Taylor, M. H. (1978). Histopathological and physiological responses of Fundulus heteroclitus to naphthalene exposure. Journal of the Fisheries Research Board of Canada, 35, 1060–1066. https://doi.org/ 10.1139/f78-169
- Dorval, J., Leblond, V. S., & Hontela, A. (2003). Oxidative stress and loss of cortisol secretion in adrenocortical cells of rainbow trout (*Oncorhynchus mykiss*) exposed in vitro to endosulfan, an organochlorine pesticide. Aquatic Toxicology, 63, 229–241. https://doi.org/10.1016/S0166-445X (02)00182-0
- dos Santos, T. C. A., Ngan, P. V., de Arruda Campos Rocha Passos, M. J., & Gomes, V. (2006). Effects of naphthalene on metabolic rate and ammonia excretion of juvenile Florida pompano, *Trachinotus carolinus*.

Journal of Experimental Marine Biology and Ecology, 335, 82–90. https://doi.org/10.1016/j.jembe.2006.02.019

- Dupuis, A., & Ucan-Marin, F. (2015). A literature review on the aquatic toxicology of petroleum oil: An overview of oil properties and effects to aquatic biota (Res. Doc. 2015/007). Canadian Science Advisory Secretariat. http://www.dfo-mpo.gc.ca/csas-sccs/Publications/ResDocs-DocRech/2015/2015_007-eng.html
- Environment and Climate Change Canada, Fisheries and Oceans Canada, & Natural Resources Canada. (2013). Federal government technical report: Properties, composition and marine spill behaviour, fate and transport of two diluted bitumen products from the Canadian oil sands. http://publications.gc.ca/collections/collection_2014/ec/En84-96-2013eng.pdf
- Farrell, A. P. (2008). Comparisons of swimming performance in rainbow trout using constant acceleration and critical swimming speed tests. *Journal of Fish Biology*, 72, 693–710. https://doi.org/10.1111/j.1095-8649.2007.01759.x
- Folch, J., Lees, M., & Stanley, G. H. S. (1957). A simple method for the isolation and purification of total lipids from animal tissues. *Journal of Biological Chemistry*, 226, 497–509.
- Gregory, T. R., & Wood, C. M. (1999). The effects of chronic plasma cortisol elevation on the feeding behaviour, growth, competitive ability, and swimming performance of juvenile rainbow trout. *Physiological and Bi*ochemical Zoology, 72, 286–295. https://doi.org/10.1086/316673
- Hammer, C. (1995). Fatigue and exercise tests with fish. Comparative Biochemistry and Physiology Part A: Physiology, 112, 1–20. https://doi.org/ 10.1016/0300-9629(95)00060-K
- Heintz, R. A., Short, J. W., & Rice, S. D. (1999). Sensitivity of fish embryos to weathered crude oil: Part II. Increased mortality of pink salmon (Oncorhynchus gorbuscha) embryos incubating downstream from weathered Exxon Valdez crude oil. Environmental Toxicology and Chemistry, 18, 494–503. https://doi.org/10.1002/etc.5620180318
- Henderson, M. A., & Graham, C. (1998). History and status of Pacific salmon in British Columbia. North Pacific Anadromous Fish Commission Bulletin, 1, 13–22.
- Hicken, C. E., Linbo, T. L., Baldwin, D. H., Willis, M. L., Myers, M. S., Holland, L., Larsen, M., Stekoll, M. S., Rice, S. D., Collier, T. K., Scholz, N. L., & Incardona, J. P. (2011). Sublethal exposure to crude oil during embryonic development alters cardiac morphology and reduces aerobic capacity in adult fish. Proceedings of the National Academy of Sciences of the United States of America, 108, 7086–7090. https://doi.org/10.1073/ pnas.1019031108
- Hontela, A. (1997). Endocrine and physiological responses of fish to xenobiotics: Role of glucocorticosteroid hormones. *Reviews in Toxicology*, 1, 159–206.
- Incardona, J. P. (2017). Molecular mechanisms of crude oil developmental toxicity in fish. Archives of Environmental Contamination and Toxicology, 73, 19–32. https://doi.org/10.1007/s00244-017-0381-1
- Johansen, J. L., & Esbaugh, A. J. (2017). Sustained impairment of respiratory function and swim performance following acute oil exposure in a coastal marine fish. Aquatic Toxicology, 187, 82–89. https://doi.org/10.1016/j. aquatox.2017.04.002
- Kennedy, C. J. (2014). Multiple effects of oil and its components in fish. In J. Alford, M. Peterson, & C. Green (Eds.), Impacts of oil spill disasters on marine habitats and fisheries in North America (pp. 3–34). CRC.
- Kennedy, C. J., & Farrell, A. P. (2005). Ion homeostasis and interrenal stress responses in juvenile Pacific herring, *Clupea pallasi*, exposed to the water-soluble fraction of crude oil. *Journal of Experimental Marine Biology and Ecology*, 323, 43–56. https://doi.org/10.1016/j.jembe.2005. 02.021
- Kennedy, C. J., & Farrell, A. P. (2006). Effects of exposure to the watersoluble fraction of crude oil on the swimming performance and the metabolic and ionic recovery postexercise in Pacific herring (*Clupea pallasi*). Environmental Toxicology and Chemistry, 25, 2715–2724. https://doi.org/10.1897/05-504R.1
- Klinger, D. H., Dale, J. J., Machado, B. E., Incardona, J. P., Farwell, C. J., & Block, B. A. (2015). Exposure to Deepwater Horizon weathered crude oil increases routine metabolic demand in chub mackerel, *Scomber japonicus*. *Marine Pollution Bulletin*, *98*, 259–266. https://doi.org/10.1016/j. marpolbul.2015.06.039
- Kochhann, D., Meyersieck Jardim, M., Valdez Domingos, F. X., & Luis Val, A. (2015). Biochemical and behavioral responses of the Amazonian fish *Colossoma macropomum* to crude oil: The effect of oil layer on water

surface. Ecotoxicology and Environmental Safety, 111, 32–41. https://doi.org/10.1016/j.ecoenv.2014.09.016

- Labelle, M. (2009). Status of Pacific salmon resources in southern British Columbia and the Fraser River basin. Pacific Fisheries Resource Conservation Council.
- Laurel, B. J., Copeman, L. A., Iseri, P., Spencer, M. L., Hutchinson, G., Nordtug, T., Donald, C. E., Meier, S., Allan, S. E., Boyd, D. T., Ylitalo, G. M., Cameron, J. R., French, B. L., Linbo, T. L., Scholz, N. L., & Incardona, J. P. (2019). Embryonic crude oil exposure impairs growth and lipid allocation in a keystone arctic forage fish. *iScience*, 19, 1101–1113. https://doi.org/10.1016/j.isci.2019.08.051
- Levy, D. A. (2009). Pipelines and salmon in northern British Columbia: Potential impacts. Pembina Institute. https://www.pembinainstitute.org/ reports/pipelines-and-salmon-in-northern-bc-report.pdf
- Lin, F., Baillon, L., Langlois, V. S., & Kennedy, C. J. (2021). Environmental modulators of diluted bitumen effects in juvenile pink salmon (Oncorhynchus gorbuscha). Marine Environmental Research, 169, Article 105392. https://doi.org/10.1016/j.marenvres.2021.105392
- Lin, F., Osachoff, H. L., & Kennedy, C. J. (2020). Physiological disturbances in juvenile sockeye salmon (*Oncorhynchus nerka*) exposed to the watersoluble fraction of diluted bitumen. *Aquatic Toxicology*, 220, Article 105383. https://doi.org/10.1016/j.aquatox.2019.105383
- Lockhart, W. L., Duncan, D. A., Billeck, B. N., Danell, R. A., & Ryan, M. J. (1996). Chronic toxicity of the "water-soluble fraction" of Norman Wells crude oil to juvenile fish. Spill Science and Technology Bulletin, 3, 259–262. https://doi.org/10.1016/S1353-2561(97)00024-8
- López-Patiño, M. A., Hernández-Pérez, J., Gesto, M., Librán-Pérez, M., Míguez, J. M., & Soengas, J. L. (2014). Short-term time course of liver metabolic response to acute handling stress in rainbow trout, Oncorhynchus mykiss. Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology, 168, 40–49. https://doi.org/10.1016/ j.cbpa.2013.10.027
- Madison, B. N., Hodson, P. V., & Langlois, V. S. (2015). Diluted bitumen causes deformities and molecular responses indicative of oxidative stress in Japanese medaka embryos. *Aquatic Toxicology*, *165*, 222–230. https://doi.org/10.1016/j.aquatox.2015.06.006
- Madison, B. N., Hodson, P. V., & Langlois, V. S. (2017). Cold Lake Blend diluted bitumen toxicity to the early development of Japanese medaka. *Environmental Pollution*, 225, 579–586. https://doi.org/10.1016/j.envpol. 2017.03.025
- Madison, B. N., Tavakoli, S., Kramer, S., & Bernier, N. J. (2015). Chronic cortisol and the regulation of food intake and the endocrine growth axis in rainbow trout. *Journal of Endocrinology*, 226, 103–119. https://doi. org/10.1530/JOE-15-0186
- Mager, E. M., Esbaugh, A. J., Stieglitz, J. D., Hoenig, R., Bodinier, C., Incardona, J. P., Scholz, N. L., Benetti, D. D., & Grosell, M. (2014). Acute embryonic or juvenile exposure to Deepwater Horizon crude oil impairs the swimming performance of mahi-mahi (*Coryphaena hippurus*). *Envi*ronmental Science & Technology, 48, 7053–7061. https://doi.org/10. 1021/es501628k
- McDonnell, D., Madison, B. N., Baillon, L., Wallace, S. J., Brown, S. R., Hodson, P. V., & Langlois, V. S. (2019). Comparative toxicity of two diluted bitumens to developing yellow perch (*Perca flavescens*). *Science* of the Total Environment, 655, 977–985. https://doi.org/10.1016/j. scitotenv.2018.11.199
- McPhee, D. L., & Janz, D. M. (2014). Dietary selenomethionine exposure alters swimming performance, metabolic capacity and energy homeostasis in juvenile fathead minnow. *Aquatic Toxicology*, 155, 91–100. https://doi.org/10.1016/j.aquatox.2014.06.012
- Meador, J. P., Sommers, F. C., Ylitalo, G. M., & Sloan, C. A. (2006). Altered growth and related physiological responses in juvenile Chinook salmon (Oncorhynchus tshawytscha) from dietary exposure to polycyclic aromatic hydrocarbons (PAHs). Canadian Journal of Fisheries and Aquatic Sciences, 63, 2364–2376. https://doi.org/10.1139/f06-127
- Milligan, C. L. (2003). A regulatory role for cortisol in muscle glycogen metabolism in rainbow trout Oncorhynchus mykiss Walbaum. Journal of Experimental Biology, 206, 3167–3173. https://doi.org/10.1242/jeb.00538
- Moles, A., & Rice, S. D. (1983). Effects of crude oil and naphthalene on growth, caloric content, and fat content of pink salmon juveniles in seawater. *Transactions of the American Fisheries Society*, 112, 205–211. https://doi.org/10.1577/1548-8659(1983)112<205:EOCOAN>2.0.CO;2
- Mommsen, T. P., Vijayan, M. M., & Moon, T. W. (1999). Cortisol in teleosts: Dynamics, mechanisms of action, and metabolic regulation. *Reviews in*

Fish Biology and Fisheries, 9, 211–268. https://doi.org/10.1023/ A:1008924418720

- Moyes, C. D., & West, T. G. (1995). Exercise metabolism of fish. In P. W. Hochachka & T. P. Mommsen (Eds.), *Biochemistry and molecular biology* of fishes: Vol. 4. Metabolic biochemistry (pp. 367–392). Elsevier. https:// doi.org/10.1016/S1873-0140(06)80019-6
- National Academies of Sciences, Engineering, and Medicine. (2016). Spills of diluted bitumen from pipelines: A comparative study of environmental fate, effects, and response. National Academies.
- National Energy Board. (2019). Optimizing oil pipeline and rail capacity out of western Canada—Advice to the minister of natural resources. https:// www.cer-rec.gc.ca/nrg/sttstc/crdIndptrImprdct/rprt/2019ptmzngcpct/ index-eng.html
- Natural Resources Canada. (2020). Crude oil industry overview. https://www. nrcan.gc.ca/our-natural-resources/energy-sources-distribution/fossilfuels/crude-oil/crude-oil-industry-overview/18078
- Nelson, D., Stieglitz, J. D., Cox, G. K., Heuer, R. M., Benetti, D. D., Grosell, M., & Crossley, D. A. (2017). Cardio-respiratory function during exercise in the cobia, *Rachycentron canadum*: The impact of crude oil exposure. *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology*, 201, 58–65. https://doi.org/10.1016/j.cbpc.2017.08.006
- Nendick, L., Grant, A., Gardner, M., Sackville, M., Brauner, C. J., & Farrell, A. P. (2009). Swimming performance and associated ionic disturbance of juvenile pink salmon Oncorhynchus gorbuscha determined using different acceleration profiles. Journal of Fish Biology, 75, 1626–1638. https://doi.org/10.1111/j.1095-8649.2009.02388.x
- Ontario Ministry of Natural Resources. (2009). Egg disinfection and incubation procedures for salmonids (salmon, trout, and whitefish). *Fish Culture Technical Bulletin*, 1, 1–9. Retrieved November 2019, from: https://dr6j45jk9xcmk.cloudfront.net/documents/2545/268425.pdf
- Osachoff, H. L., Osachoff, K. N., Wickramaratne, A. E., Gunawardane, E. K., Venturini, F. P., & Kennedy, C. J. (2014). Altered burst swimming in rainbow trout Oncorhynchus mykiss exposed to natural and synthetic oestrogens. Journal of Fish Biology, 85, 210–227. https://doi.org/10. 1111/jfb.12403
- Pasparakis, C., Mager, E. M., Stieglitz, J. D., Benetti, D., & Grosell, M. (2016). Effects of Deepwater Horizon crude oil exposure, temperature and developmental stage on oxygen consumption of embryonic and larval mahi-mahi (*Coryphaena hippurus*). Aquatic Toxicology, 181, 113–123. https://doi.org/10.1016/j.aquatox.2016.10.022
- Philibert, D. A., Philibert, C. P., Lewis, C., & Tierney, K. B. (2016). Comparison of diluted bitumen (dilbit) and conventional crude oil toxicity to developing zebrafish. *Environmental Science & Technology*, 50, 6091–6098. https://doi.org/10.1021/acs.est.6b00949
- Raincoast Conservation Foundation. (2018). Executive summary: Wild salmon, pipelines, and the Trans Mountain expansion. https://www.raincoast.org/reports/salmon-oil-pipeline/
- Sørhus, E., Incardona, J. P., Furmanek, T., Goetz, G. W., Scholz, N. L., Meier, S., Edvardsen, R. B., & Jentoft, S. (2017). Novel adverse outcome pathways revealed by chemical genetics in a developing marine fish. *eLife*, 6, Article e20707. https://doi.org/10.7554/eLife.20707
- Srivastava, R. K., & Brown, J. A. (1991). The biochemical characteristics and hatching performance of cultured and wild Atlantic salmon (*Salmo salar*) eggs. *Canadian Journal of Zoology*, 69, 2436–2441. https://doi.org/10. 1139/z91-342

- Stieglitz, J. D., Mager, E. M., Hoenig, R. H., Benetti, D. D., & Grosell, M. (2016). Impacts of Deepwater Horizon crude oil exposure on adult mahimahi (Coryphaena hippurus) swim performance. Environmental Toxicology and Chemistry, 35, 2613–2622. https://doi.org/10.1002/etc.3436
- Thomas, P., Woodin, B. R., & Neff, J. M. (1980). Biochemical responses of the striped mullet *Mugil cephalus* to oil exposure I. Acute responses—Interrenal activations and secondary stress responses. *Marine Biology*, 59, 141–149. https://doi.org/10.1007/ BF00396861
- Thomas, R. E., & Rice, S. D. (1987). Effect of water-soluble fraction of cook inlet crude oil on swimming performance and plasma cortisol in juvenile coho salmon (Oncorhynchus kisutch). Comparative Biochemistry and Physiology Part C: Comparative Pharmacology, 87, 177–180. https://doi. org/10.1016/0742-8413(87)90200-3
- US Environmental Protection Agency. (2013). Dredging begins on Kalamazoo River: Enbridge oil spill, Marshall, Michigan. https://www. epa.gov/enbridge-spill-michigan/enbridge-oil-spill-fact-sheets
- Vignier, V., Vandermeulen, J. H., & Fraser, A. J. (1992). Growth and food conversion by Atlantic salmon parr during 40 days' exposure to crude oil. *Transactions of the American Fisheries Society*, 121, 322–332. https://doi.org/10.1577/1548-8659(1992)121<0322:GAFCBA>2.3.CO;2
- Vijayan, M. M., Ballantyne, J. S., & Leatherland, J. F. (1991). Cortisolinduced changes in some aspects of the intermediary metabolism of Salvelinus fontinalis. General and Comparative Endocrinology, 82, 476–486. https://doi.org/10.1016/0016-6480(91)90323-X
- Wang, S. Y., Lum, J. L., Carls, M. G., & Rice, S. D. (1993). Relationship between growth and total nucleic acids in juvenile pink salmon, Oncorhynchus gorbuscha, fed crude oil contaminated food. Canadian Journal of Fisheries and Aquatic Sciences, 50, 996–1001. https://doi.org/10. 1139/f93-115
- Weber, L. P., Dubé, M. G., Rickwood, C. J., Driedger, K., Portt, C., Brereton, C., & Janz, D. M. (2008). Effects of multiple effluents on resident fish from Junction Creek, Sudbury, Ontario. *Ecotoxicology and Environmental Safety*, 70, 433–445. https://doi.org/10.1016/j.ecoenv.2007. 08.001
- Weber, L. P., Higgins, P. S., Carlson, R. I., & Janz, D. M. (2003). Development and validation of methods for measuring multiple biochemical indices of condition in juvenile fishes. *Journal of Fish Biology*, 63, 637–658. https://doi.org/10.1046/j.1095-8649.2003.00178.x
- Woodward, D. F., Riley, R. G., & Smith, C. E. (1983). Accumulation, sublethal effects, and safe concentration of a refined oil as evaluated with cutthroat trout. Archives of Environmental Contamination and Toxicology, 12, 455–464. https://doi.org/10.1007/BF01057589
- Xu, E. G., Khursigara, A. J., Magnuson, J., Hazard, E. S., Hardiman, G., Esbaugh, A. J., Roberts, A. P., & Schlenk, D. (2017). Larval red drum (*Sciaenops ocellatus*) sublethal exposure to weathered Deepwater Horizon crude oil: Developmental and transcriptomic consequences. *Environmental Science & Technology*, *51*, 10162–10172. https://doi.org/ 10.1021/acs.est.7b02037
- Xu, E. G., Mager, E. M., Grosell, M., Pasparakis, C., Schlenker, L. S., Stieglitz, J. D., Benetti, D., Hazard, E. S., Courtney, S. M., Diamante, G., Freitas, J., Hardiman, G., & Schlenk, D. (2016). Time- and oil-dependent transcriptomic and physiological responses to Deepwater Horizon oil in mahi-mahi (*Coryphaena hippurus*) embryos and larvae. *Environmental Science & Technology*, *50*, 7842–7851. https://doi.org/10.1021/acs.est. 6b02205